

Reversible PO Nanofiber Membrane Chromatography Syringe Capture Viruses And Viral Vectors 13mm 0.70μM

Our Product Introduction

for more products please visit us on wechanfib.com

Basic Information

- Place of Origin: PRC
- Brand Name: WECHANFIB
- Model Number: WF-ZT-ST-MCS-13-70
- Minimum Order Quantity: 100
- Price: Negotiable
- Packaging Details: Individually blister package, 50-100per box
- Delivery Time: 24-30days
- Payment Terms: L/C, D/A, D/P, T/T, Western Union, MoneyGram, Paypal
- Supply Ability: 5000pc/month



Product Specification

- Membrane Material: EVOH
- Shell Material: PP
- Diameter: 13mm
- Pore Size: 0.70μm
- Sterile: Yes
- Application Type: MCQ Anion
- Highlight:
 - PO Nanofiber Membrane Chromatography Syringe Capture
 - 13mm Nanofiber Membrane Chromatography Syringe Capture
 - 0.70μm Nanofiber Membrane Chromatography Syringe Capture



Product Description

PO Nanofiber Membrane Chromatography Syringe Caputer viruses and viral vectors 13mm 0.70μm MCQ Anion Ion Exchange Chromatography

Description MCQ Anion Ion Exchange Chromatography

Ion exchange chromatography is a technique for separating compounds based on their net charge. Ion exchange chromatography media contain negatively or positively charged functional groups covalently bound to a solid support, yielding either a cation or anion exchanger, respectively. Charged compounds are adsorbed and retained by an ion exchanger having the opposite charge, whereas compounds that are neutral or have the same charge as the media pass through the void volume and are eluted from the column. The binding of the charged compounds is reversible, and adsorbed compounds are commonly eluted with a salt or pH gradient. Ion exchange media are available in various particle sizes, ionic forms, and purity ranges.

Application MCQ Anion Ion Exchange Chromatography

Capture large particles such as recombinant proteins, plasmids, viruses, viral vectors and plasma components; Polishing steps to remove contaminants such as host DNA, HCPs, and endotoxins in biological fluids; Fine purification of small molecule drugs such as oligonucleotides and peptides.

Features MCQ Anion Ion Exchange Chromatography

Materials to clarify both aqueous and mild organic solutions;
100% passed integrity test;
Low-binding membrane;
Low IC extractables;
Ethylene oxide or gamma irradiation sterilization;

Our Product Introduction

Ready to use equipment;
Easy to connect Syringe Filter or Pump;
High adsorption capacity to macromolecule;
High flow rate and high capacity;
Well developed separation function.

Filter Performance MCQ Anion Ion Exchange Chromatography

Type	Anion	Cation
Membrane Materials	PO	PO
Shell Materials	PP	PP
Membrane Area	20cm ²	20cm ²
Filter Diameter	25mm	25mm
Ligand Type(Density)	Amino(4mmol/g)*	Carboxyl(2.6mmol/g)
Dynamic Loading	136mg/g(RNA)**	510mg/g(Lysozyme)
Pore Size	0.7µm	0.7µm
Porosity	78%	78%
Flow Rate	0.1-100ml/min	0.1-100ml/min
Recommended Flow Rate	1ml/min	1ml/min
Operating Pressure	1-10bar	1-10bar
Reusability	>10cycles	>10cycles
Package	Individual Package	Individual Package
Product Code	WF-ZT-ST-MCQ-25-70	WF-ZT-ST-MCS-25-70
Hold-up Volume	1ml	1ml
pH Range	3-14	3-14
Sample Storage Material	PP	PP
Sample Volume	200-1000µl	200-1000µl
Max. Operating Temperature	40	40
Min. Operating Temperature	0	0
Note: *Quaternary ammonium salt ligand products are also available **DNA and BSA data is available.		

Naming Regulations MCQ Anion Ion Exchange Chromatography

WF	ZT	ST	SWG(式无)	MCQ	25	70	2.5(式无)
Trademark	Type	Sterilize	Connector	Application	Diameter	Pore Size	Length
WECHANFIB	ZT Syringe	Sterilize	SWG Intubation	MCS Cation	13mm	45 0.45µm	2.5 2.5"
	NL Capsule		MNPT Whorl	MCQ Anion	25mm	70 0.70µm	5 5"
	CT Centrifuge tube			MCPA Protein A	33mm		10 10"
							20 20"

维晨科技有限公司

WECHANGE TECHNOLOGY CO.,LTD

We-Change Makes Your Life better

(1) NFMC Ion Exchange Chromatography Membrane

WECHANFIB NFMC Ion Exchange Chromatography Membrane uses stable porous structure PO(EVOH) nanofiber membrane, its over 0.7µm pore size which is much more greater than normal gel chromatography medium. Entrained by the flow, molecules have been carried to the functional groups fixed on the surface of ion exchange membranes rapidly. The adsorption capacity of ion exchange ligands is relatively stable and will not loss after repeated use. Our NFMC ion exchange membranes have strong/weak anions(Q/PA) & strong cation(S) types, effective area ranging from 0.35-100cm², corresponding to adsorb proteins 10-100mg per unit. They can be used together with our syringe filters and centrifugal purification columns, our syringe filter equipped with standard Luer-Lok fitting, can be connected with HPLC or FPLC system, it is compatible with various solvents, repeat use for more than 100 times. Our centrifugal purification columns help you to achieve separation in 5 min.



Type	Anion	Cation
Membrane Materials	PO	PO
Shell Materials	PP	PP
Membrane Area	20cm ²	20cm ²
Filter Diameter	25mm	25mm
Ligand Type (Density)	Amino(4mmol/g)*	Carboxyl(2.6mmol/g)
Dynamic Loading	136mg/g(RNA)**	510mg/g(Lysozyme)
Pore Size	0.7µm	0.7µm
Porosity	78%	78%
Flow Rate	0.1-100ml/min	0.1-100ml/min
Recommended Flow Rate	1ml/min	1ml/min
Operating Pressure	1-10bar	1-10bar
Reusability	>10cycles	>10cycles
Package	Individual Package	Individual Package
Product Code	WF-ZT-ST-MCQ-25-70	WF-ZT-ST-MCS-25-70

Type	Cation
Effective Filtration Area	0.35cm ²
Membrane Materials	PO
Hold-up Volume	1ml
Maximum Centrifugal Force	14000g
pH Range Maximum	14
pH Range Minimum	3
Sample Storage Material	PP
Sample Volume	200-1000µl
Maximum Operating Temperature	40°C
Minimum Operating Temperature	0°C
Pore Size	700nm
Product Code	WF-CT-ST-MCQ-35-70

Note: * Quaternary ammonium salt ligand products are also available.
** DNA and BSA data is available.

WECHANFIB
维晨飞昂

Wuhan Wechange Technology Co., Ltd



sales@wechange.com.cn



wechanfib.com

Room 506, 5th Floor, Block B, Kechuang Plaza, No. 10, Wenhua Avenue, Jiangxia District, Wuhan, Hubei Province, China